Table 44. Maternal mortality for complications of pregnancy, childbirth, and the puerperium, according to race, Hispanic origin, and age: United States, selected years 1950–2000

[Data are based on death certificates]

Race, Hispanic origin, and age	1950¹	1960¹	1970	1980	1990	1995	1997	1998	1999²	2000²
	Number of deaths									
All persons	2,960	1,579	803	334	343	277	327	281	391	396
White	1,873 1,041 	936 624 	445 342 	193 127 3 11	177 153 4 9	129 133 1 14	179 125 2 21	158 104 2 17	214 154 5 18	240 137 6 13
Hispanic <sup>3</sup>					47 125	43 84	57 121	42 116	67 149	81 160
All persons	Deaths per 100,000 live births									
All ages, age adjusted	73.7 83.3	32.1 37.1	21.5 21.5	9.4 9.2	7.6 8.2	6.3 7.1	7.6 8.4	6.1 7.1	8.3 9.9	8.2 9.8
Under 20 years	70.7 47.6 63.5 107.7 222.0	22.7 20.7 29.8 50.3 104.3	18.9 13.0 17.0 31.6 81.9	7.6 5.8 7.7 13.6 36.3	7.5 6.1 6.0 9.5 20.7	3.9 5.7 6.0 7.3 15.9	5.7 6.6 7.9 8.3 16.1	5.0 6.7 7.5 14.5	6.6 6.2 8.2 10.1 23.0	7.4 7.9 10.0 22.7
White									1	
All ages, age adjusted	53.1 61.1	22.4 26.0	14.4 14.3	6.7 6.6	5.1 5.4	3.6 4.2	5.2 5.8	4.2 5.1	5.5 6.8	6.2 7.5
Under 20 years	44.9 35.7 45.0 75.9 174.1	14.8 15.3 20.3 34.3 73.9	13.8 8.4 11.1 18.7 59.3	5.8 4.2 5.4 9.3 25.5	3.9 4.8 5.0 12.6	3.5 4.0 4.0 9.1	4.2 5.4 5.4 11.5	3.1 4.9 4.9 11.0	4.0 5.4 7.0 16.6	5.6 5.9 7.1 18.0
Black										
All ages, age adjusted		92.0 103.6	65.5 60.9	24.9 22.4	21.7 22.4	20.9 22.1	20.1 20.8	16.1 17.1	23.3 25.4	20.1 22.0
Under 20 years		54.8 56.9 92.8 150.6 299.5	32.3 41.9 65.2 117.8 207.5	13.1 13.9 22.4 44.0 100.6	14.7 14.9 44.2 79.7	15.3 21.0 31.2 61.4	15.3 24.3 32.9 40.4	* 12.7 17.2 27.7 37.2	14.0 26.6 36.1 69.9	15.3 21.8 34.8 62.8
Hispanic <sup>3,5</sup>									I	
All ages, age adjusted					7.4 7.9	5.4 6.3	7.6 8.0	5.2 5.7	7.9 8.8	9.0 9.9
White, non-Hispanic <sup>3</sup>									1	
All ages, age adjusted					4.4 4.8	3.3 3.5	4.4 5.2	4.0 4.9	4.9 6.4	5.5 6.8

<sup>- - -</sup> Data not available.

NOTES: Rates are age adjusted to the 1970 distribution of live births by mother's age in the United States. See Appendix II, Age adjustment. For data years shown, the code numbers for cause of death are based on the then current *International Classification of Diseases*, described in Appendix II, tables IV and V. The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Conversely, persons of Hispanic origin may be of any race. For 1950 and 1960, rates are based on live births by race of child; for all other years, rates are based on live births by race of mother. See Appendix I, National Vital Statistics System. Rates are not calculated for American Indian or Alaska Native and Asian or Pacific Islander mothers because rates based on fewer than 20 deaths are unreliable. Data for additional years are available (see Appendix III).

SOURCES: Centers for Disease Control and Prevention, National Center for Health Statistics, National Vital Statistics System; Vital statistics of the United States, vol I, natality and vol II, mortality, part A, for data years 1950–93. Public Health Service. Washington: U.S. Government Printing Office; for 1994–2000, unpublished data.

52 Health, United States, 2002

<sup>\*</sup> Based on fewer than 20 deaths.

<sup>&</sup>lt;sup>1</sup>Includes deaths of persons who were not residents of the 50 States and the District of Columbia.

<sup>&</sup>lt;sup>2</sup>Starting with 1999 data, changes have been made in the classification and coding of maternal deaths under ICD–10. The large increase in the number of maternal deaths between 1998 and 1999 is due to changes associated with ICD–10. See Appendix II, *International Classification of Diseases (ICD)*; and Rate: Death and related rates.

<sup>&</sup>lt;sup>3</sup>Excludes data from States lacking an Hispanic-origin item on their death and birth certificates. See Appendix I, National Vital Statistics System.

<sup>&</sup>lt;sup>4</sup>Rates computed by relating deaths of women 35 years and over to live births to women 35–49 years.

<sup>&</sup>lt;sup>5</sup>Age-specific maternal mortality rates are not calculated because rates based on fewer than 20 deaths are unreliable.